

# Astro2020 Science White Paper

## The Sun-like Stars Opportunity

**Thematic Areas:** Planetary Systems

### **Principal Author:**

**Name:** Giada N. Arney

**Institution:** NASA Goddard Space Flight Center

**Email:** giada.n.arney@nasa.gov

**Phone:** 301-614-6627

### **Co-authors:**

Natasha Batalha (UC Santa Cruz), Amber V. Britt (Fisk University), Nicolas Cowan (McGill), Shawn D. Domagal-Goldman (NASA GSFC), Courtney Dressing (UC Berkeley), Kevin France (CU Boulder), Yuka Fujii (ELSI), Ravi Kopparapu (NASA GSFC), Stephen Kane (UC Riverside), Joshua Krissansen-Totton (UW), Andrew Lincowski (UW), Owen Lehmer (UW, NASA Ames), Eric Lopez (NASA GSFC), Jacob Lustig-Yaeger (UW), Victoria S. Meadows (UW), Stephanie Olson (Chicago), M. Niki Paranteau (NASA Ames), Ilaria Pascucci (LPL), Ramses Ramirez (ELSI), Christopher Reinhard (GA Tech), Aki Roberge (NASA GSFC), Tyler D. Robinson (NAU), Edward Schwieterman (UC Riverside), Christopher Stark (STScI), Eric. T. Wolf (CU Boulder), Allison Youngblood (NASA GSFC)

### **Co-signers:**

Mahmuda Afrin Badhan (UMD), Jacob Bean (University of Chicago), Alexandra Brosius (Penn State), James B. Breckinridge (Caltech, Arizona), Douglas A. Caldwell (SETI Exoplanets), Knicole Colón (NASA GSFC), Joleen K. Carlberg (STScI), William Danchi (NASA GSFC), James R. A. Davenport (UW), Anthony Del Genio (NASA GISS), Chuanfei Dong (Princeton University), Diana Dragomir (MIT/UNM), Thomas Fauchez (NASA GSFC), Debra Fischer (Yale University), B. Scott Gaudi (Ohio State University), Julien H. Girard (STScI), David Grinspoon (Planetary Science Institute), Tyler D. Groff (NASA GSFC), Scott D. Guzewich (NASA GSFC), Nader Haghighipour (UH-IfA), Brian Hicks (NASA GSFC/UMD), Natalie Hinkel (SwRI), Jinyoung Serena Kim (University of Arizona), Erika Kohler (NASA GSFC), Yuni Lee (NASA GSFC/UMBC), Jeffrey Linsky (University of Colorado), Timothy A. Livengood (NASA GSFC/UMCP), Timothy Lyons (UC Riverside), Eric Mamajek (JPL/Caltech), Mark Marley (NASA Ames), Dimitri Mawet (Caltech/JPL), Bertrand Mennesson (JPL), Susan Mullally (STScI), John O'Meara (Keck Observatory), Daria Pidhorodetska (NASA GSFC), Peter Plavchan (George Mason University), Sam Ragland (Keck Observatory), Sarah Rugheimer (Oxford), Caleb Scharf (Columbia U.), Evgenya Shkolnik (ASU), Keivan G. Stassun (Vanderbilt University), R. Deno Stelter (UC Santa Cruz/UC Observatories), Stuart F. Taylor (Participation Worldscape), Johanna Teske (Carnegie Observatories), Margaret Turnbull (SETI Institute), Monica Vidaurri (George Mason University), Michael Way (NASA GISS), Michael L. Wong (UW), Robin Wordsworth (Harvard University), Neil T. Zimmerman (NASA GSFC)

**Abstract:**


Observations of exoplanets around Sun-like (i.e. FGK) stars will allow us to place our planet into a larger cosmic context and may offer the best chance of finding habitable, and inhabited, planets. New facilities beyond those currently planned would be required to accomplish these observations. Exoplanet community reports have emphasized the importance of observations of exoplanets in the habitable zones of Sun-like stars for over a decade. This trend continued in the recent Exoplanet Science Strategy report. We endorse the findings and recommendations published in the National Academy reports on Exoplanet Science Strategy and Astrobiology Strategy for the Search for Life in the Universe. This white paper extends and complements the material presented therein.

# 1. Introduction

The study of exoplanets is a young field, but it remains one of the most compelling and exciting areas of astronomy with popular appeal beyond the astronomical community. The path to characterizing habitable and inhabited worlds has already begun with ground- and space-based programs discovering planets in the habitable zones (HZs) of their stars. These potentially habitable planets, especially those orbiting M dwarfs, will be prime targets in coming years for spectral observations with the James Webb Space Telescope (JWST) and large ground-based facilities. However, while they are the simplest to observe, planets orbiting M dwarfs may face multiple barriers to habitability. Terrestrial planets in the HZs of Sun-like stars (i.e. FGK dwarfs) offer the best chance of discovering planets with conditions and evolutionary histories analogous to Earth’s, as well as the best opportunity to detect unambiguous biosignatures.

## A Consensus of Reports

The importance of observations of Earthlike planets around Sun-like stars has been reiterated by the exoplanet community for over a decade. The 2008 *Report of the ExoPlanet Task Force*<sup>1</sup> by the Astronomy and Astrophysics Advisory Committee (Lunine et al., 2008) called for “finding and characterization of a planet like our own, around a star like our own Sun.” This report laid out two parallel tracks for the community: 1) study targets around M dwarfs using ground and space-based assets, and 2) invest in new technologies and space-based facilities to study planets around Sun-like stars (**Figure 1**). While major progress has and will continue to be made on the “M dwarf track” with existing and planned facilities, progress in discovering and characterizing habitable worlds around Sun-like stars still requires new space-based assets.



**Figure 1:** The recommended exoplanet science strategy from the 2008 *Report of the ExoPlanet Task Force*. Track 1 (M dwarfs) is now in progress. Track 2 (F, G, K dwarfs; i.e. “Sun-like” stars) will require major new technology and facilities for significant progress. Figure credit: Figure 1 from the *Report of the ExoPlanet Task Force* (Lunine et al., 2008).

<sup>1</sup> [https://www.nsf.gov/mps/ast/aaac/exoplanet\\_task\\_force/reports/exoptf\\_final\\_report.pdf](https://www.nsf.gov/mps/ast/aaac/exoplanet_task_force/reports/exoptf_final_report.pdf)

Other community reports in the intervening years have reached similar conclusions. The 2010 *European Roadmap for Exoplanets*<sup>2</sup> (Hatzes et al., 2010) emphasized the importance of observations of exoplanets in the HZs of Sun-like stars. Similarly, the 2013 NASA Astrophysics Roadmap *Enduring Quests Daring Visions*<sup>3</sup> (Kouveliotou et al., 2013) stated that observations of exoplanets around Sun-like stars will give us “greater hopes of identifying and understanding life” compared to planets around lower mass stars. The 2014 Exo-C and Exo-S reports on probe-scale space missions reiterated the importance of observations of exoplanets in the HZs of Sun-like stars (Stapelfeldt et al., 2014; Seager et al., 2014). Most recently, the 2018 *Exoplanet Science Strategy* report (NAS 2018) repeated this emphasis in its first recommendation: “NASA should lead a large strategic direct imaging mission capable of measuring the reflected-light spectra of temperate terrestrial planets orbiting Sun-like stars.”

## 2. Why Sun-like Stars?

Studies of planets around Sun-like stars will place our planet into a larger cosmic context and reveal how typical our world is. Furthermore, while the challenges of observing planets around Sun-like stars are not trivial, planets around these stars represent our best chance of finding life elsewhere, in the event that M dwarfs are inhospitable hosts for habitable planets. Here we will briefly outline why observations of rocky exoplanets in the HZs of FGK host stars should be sought as a high priority. Many of the reasons why Sun-like stars are important have been described in the recent *Exoplanet Science Strategy* report (NAS 2018); we concur with this report’s findings, and this white paper extends and complements the material presented there.

### The Sun-like Star Habitability Advantage

The extreme evolution and activity of M dwarf stars suggests Sun-like stars may be more hospitable hosts for habitable worlds. M dwarfs undergo an extended super-luminous (1-2 orders of magnitude brighter) pre-main sequence (PMS) phase that can last for to up to  $10^9$  years. This can cause extreme early water loss, such that planets even in the HZs of M dwarfs may be desiccated Venus-like worlds (Luger & Barnes 2015; Ramirez & Kaltenegger 2014; Tian & Ida 2015; Lincowski et al., 2018). Possibly, this fate can be avoided by late volatile delivery (Morbidelli et al., 2000; Wang & Becker 2013) or late migration of volatile-rich planets into the HZ (Luger et al., 2015). Furthermore, many M dwarfs maintain high stellar activity levels into old age (West et al., 2015; France et al., 2018). Their high EUV and X-ray luminosities (Shkolnik et al., 2014) and frequent energetic flaring (e.g. MacGregor et al., 2018; Loyd et al., 2018) may lead to severe – perhaps total – atmospheric loss on rocky HZ planets (e.g. Owen & Mohanty 2016; Airapetian et al., 2017; Dong et al., 2017; Garcia-Sage et al., 2017). However, the planets of ultra-cool dwarfs could perhaps retain sufficient water to remain habitable depending on initial water content (Bolmont et al., 2017).

---

<sup>2</sup> <http://sci.esa.int/science-e/www/object/doc.cfm?fobjectid=47854>

<sup>3</sup> <https://science.nasa.gov/science-committee/subcommittees/nac-astrophysics-subcommittee/astrophysics-roadmap>


By contrast, planets orbiting Sun-like stars will experience more moderate stellar evolution and activity. Early K dwarfs have been highlighted as especially promising hosts for habitable planets (Cuntz & Guinan 2016). Richey-Yowell et al., (2019) found that K dwarf FUV and X-ray radiation decreases after  $\sim 100$  Myr, compared to M dwarfs whose high-energy fluxes may remain constant until  $\sim 650$  Myr (Shkolnick & Barman 2014; Schneider & Shkolnick 2018). Additionally, Richey-Yowell et al., (2019) found that incident UV and X-ray fluxes are 5 - 50 times lower for planets orbiting K dwarfs in early stages of evolution compared to early-M dwarfs and 50 - 1000 times lower compared to late-M stars. This is due to less intrinsic activity from K dwarfs and the larger star-planet separations for planets in their HZs.

## The Sun-like Star Biosignature Advantage

It may be easier to interpret biosignatures (remotely observable signs of life) on exoplanets orbiting Sun-like stars compared to M dwarfs. Studies over the last decade suggest that HZ planets around M dwarfs are especially prone to developing abiotic  $O_2$  and its photochemical byproduct  $O_3$ , two of the most important biosignatures in Earth's atmosphere (e.g. Domagal-Goldman et al., 2014; Tian et al., 2014; Gao et al., 2015; Harman et al., 2015; Luger & Barnes, 2015). A summary of the four most obvious/important false positive mechanisms appears in **Figure 2**. As can be seen, only one applies to planets around any type of star (Wordsworth & Pierrehumbert, 2014); the others are expected to be applicable to M dwarf planets. The processes that lead to these abiotic "false positive" oxygen species involve water loss and photochemical effects.

False positive  $O_2$  or  $O_3$  can be effectively ruled out by the presence of reduced gases. The "classic" biosignature disequilibrium pairing for modern Earth is the simultaneous presence of  $O_2$  and methane ( $CH_4$ ), whose atmospheric abundances are orders of magnitude away from equilibrium values (Lovelock 1965, Hitchcock & Lovelock 1967; Krissansen-Totton et al., 2016). However, this classic pairing has not always been detectable throughout Earth's inhabited history, due to low oxygen abundances at early times with enhanced  $CH_4$ , and lower  $CH_4$  abundances after the rise of oxygen (Olson et al., 2016; Reinhard et al., 2017, Schwieterman et al., 2018). Interestingly, M dwarfs and mid- and late-K dwarfs appear to offer longer  $CH_4$  photochemical lifetimes in  $O_2$ -rich atmospheres, enhancing the simultaneous detectability of these gases (Segura et al., 2005; Arney 2019). However, since Sun-like stars appear to have less chance of presenting abiotic  $O_2$  to begin with, simultaneous detection of  $O_2$  and  $CH_4$  to establish biogenicity of these gases is more critical for M dwarfs than it is for Sun-like stars.

Recent work also suggests that the productivity of oxygenic photosynthesis may be limited by light availability for planets orbiting M dwarf stars (Lehmer et al., 2018). In this case, atmospheric  $O_2$  may not accumulate to detectable levels. For instance, an Earth-like planet in the HZ of a TRAPPIST-1-like star may be unable to support a biosphere as robust as modern Earth's due to a reduced incident flux of photons with sufficient energy (400-700nm) to drive the photosynthetic reaction, which is principally responsible for fueling our planet's biosphere. The biospheres on M dwarf planets may therefore be much smaller than Earth's. In this situation, the geologic processes of Earth-like planets, such as volcanic outgassing, could overwhelm any potential atmospheric biosignature (Lehmer et al., 2018). Furthermore, complex life may be more likely to evolve on planets orbiting G dwarfs than lower mass stars, since the



**Figure 2:** Possible O<sub>2</sub> and O<sub>3</sub> biosignature false positives. Top panels show atmospheric constituents. Circled gases show false positive indicators. Bottom panels indicate species that should be absent for a given false positive to apply. Only one scenario plausibly applies to Sun-like stars for planets with low non-condensable gas inventories (i.e. low-pressure atmospheres), but such a planet could be identified given constraints on total atmospheric pressure. This may be possible with sufficient signal-to-noise observations (Feng et al. 2018; discussion in T. Robinson whitepaper). Figure credit: Meadows (2017).

latter can have greater abundances of photochemical CO in their atmospheres (Schwieterman et al., 2019). CO is toxic to aerobic organisms, and high concentrations might limit the development of complex life throughout the entire HZs of late M dwarfs.

## M Dwarfs as a Necessary Complement

The argument for observations of Sun-like stars is not an argument for ignoring M dwarfs. While there are significant questions about the habitability of rocky planets in the HZs of M dwarfs, they will provide fundamental tests of our theories of planetary evolution and biology that are derived from the Earth-Sun pairing, and they may inform us of different kinds of atmospheric processes (e.g. Yang et al., 2013; Fujii et al., 2017; Kopparapu et al., 2017; Ramirez and Kaltenegger 2018; Lincowski et al., 2018). If we find that some M dwarf HZ planets are truly habitable, this means habitable conditions are robust even in the face of significant challenges. This would have profound implications for the distribution and frequency of habitable worlds and the emergence of life in the galaxy, given that M dwarfs comprise 75% of all stars.

Characterization of small planets in the HZs of M dwarf stars will be dramatically advanced with the James Webb Space Telescope and future ground-based extremely large telescopes (ELTs). Space-based observatories are required for studies of Earth-like planets sufficiently small to be rocky ( $R < 1.6$  Earth radii; Rogers 2015) in the HZs of Sun-like stars (e.g. see C. Stark's exo-Earth yield whitepaper); possibly, ELT facilities could provide complementary imaging at thermal

wavelengths for a small number of targets, although the most detectable worlds around Sun-like stars from the ground may be too hot, or too large to be habitable (Quanz et al., 2015). An observatory that can directly image Earth-like planets around Sun-like stars would also enable high-fidelity transit spectroscopy of exoplanets around M dwarfs. **Table 1** summarizes challenges and opportunities of observations of exoplanets orbiting M dwarfs vs. FGK dwarfs.

### 3. Conclusions

The most complete understanding of the nature and distribution of habitable environments and life in the universe demands observations of multiple diverse planets orbiting stars with a range of masses. However, M dwarf worlds, although numerous and observable with upcoming facilities like JWST and ground-based ELTs, may not be habitable. To place our planet into a larger cosmic context and maximize our chances of detecting life elsewhere, observations of potentially habitable exoplanets orbiting Sun-like stars are needed. Such observations will require direct imaging using a next generation space-based telescope equipped with advanced starlight suppression technology.

**Table 1:** Challenges and opportunities for HZ planets around M dwarfs and FGK dwarfs.

	Challenges	Opportunities
<b>Potentially habitable planets orbiting M dwarfs</b>	<p>Multiple barriers to habitability possible; e.g. desiccation during super-luminous pre-main sequence phase, stellar activity-driven atmospheric loss.</p> <p>Apparent high potential for false positive oxygen biosignatures.</p> <p>Challenging to observe via direct imaging due to the small planet-star angular separation. Even nearest planets require ELTs (Crossfield 2013).</p>	<p>Most common type of stars (~75%).</p> <p>Planets with significantly different histories to Earth valuable for comparative planetology.</p> <p>Transits of HZ planets more easily observed due to shorter orbital periods and deeper transit depths. May be possible with JWST and/or ELTs.</p> <p>Moderate planet-star contrast ratio (<math>10^{-8}</math> in reflected light). Nearest planets may be within reach of ELTs (Crossfield 2013).</p>
<b>Potentially habitable planets orbiting FGK dwarfs</b>	<p>Less common types of stars.</p> <p>Transit spectroscopy prohibitively difficult due to long orbital periods, small transit depths, lower transit probability.</p> <p>Challenging planet-star contrast ratio (<math>10^{-10}</math> for a Sun-Earth twin in reflected light) demands starlight suppression advances.</p>	<p>Less significant barriers to habitability due to moderate stellar evolution and activity.</p> <p>Understand Earth in the context of planets with similar histories. Is Earth typical?</p> <p>Observable with potential future space-based telescopes in direct imaging due to larger planet-star angular separation.</p> <p>Known that life can flourish on such planets. May be best opportunity for high-confidence detection of biosignatures.</p>

## References

- Arney, G. (2019) The K Dwarf Advantage for Biosignatures on Directly Imaged Exoplanets. *The Astrophysical Journal Letters*, 873(1). <https://doi.org/10.3847/2041-8213/ab0651>
- Airapetian, V. S., et al., (2017). How hospitable are space weather affected habitable zones? The role of ion escape. *The Astrophysical Journal Letters*, 836(1), L3. <https://doi.org/10.3847/2041-8213/836/1/L3>
- Bolmont, et al. (2016). Water loss from terrestrial planets orbiting ultracool dwarfs: implications for the planets of TRAPPIST-1. *Monthly Notices of the Royal Astronomical Society*, 464(3), 3728-3741. <https://doi.org/10.1093/mnras/stw2578>
- Cuntz, M., & Guinan, E. F. (2016). About exobiology: the case for dwarf K stars. *The Astrophysical Journal*, 827(1), 79. <https://doi.org/10.3847/0004-637X/827/1/79>
- Crossfield, I. J. (2013). On high-contrast characterization of nearby, short-period exoplanets with giant segmented-mirror telescopes. *Astronomy & Astrophysics*, 551, A99. <https://doi.org/10.1051/0004-6361/201220914>
- Domagal-Goldman, S. D., Segura, A., Claire, M. W., Robinson, T. D., & Meadows, V. S. (2014). Abiotic ozone and oxygen in atmospheres similar to prebiotic Earth. *The Astrophysical Journal*, 792(2), 90. <https://doi.org/10.1088/0004-637X/792/2/90>
- Dong, C., Lingam, M., Ma, Y., & Cohen, O. (2017). Is Proxima Centauri b habitable? A study of atmospheric loss. *The Astrophysical Journal Letters*, 837(2), L26. <https://doi.org/10.3847/2041-8213/aa6438>
- France, K., et al. (2018). Far-ultraviolet Activity Levels of F, G, K, and M Dwarf Exoplanet Host Stars. *The Astrophysical Journal Supplement Series*, 239(1), 16. <https://doi.org/10.3847/1538-4365/aae1a3>
- Gao, P., Hu, R., Robinson, T. D., Li, C., & Yung, Y. L. (2015). Stability of CO<sub>2</sub> atmospheres on desiccated M dwarf exoplanets. *The Astrophysical Journal*, 806(2), 249. <https://doi.org/10.1088/0004-637X/806/2/249>



Garcia-Sage, K., Gloer, A., Drake, J. J., Gronoff, G., & Cohen, O. (2017). On the magnetic protection of the atmosphere of Proxima Centauri b. *The Astrophysical Journal Letters*, 844(1), L13. <https://doi.org/10.3847/2041-8213/aa7eca>

Fujii, Y., Del Genio, A. D., & Amundsen, D. S. (2017). NIR-driven moist upper atmospheres of synchronously rotating temperate terrestrial exoplanets. *The Astrophysical Journal*, 848(2), 100. <https://doi.org/10.3847/1538-4357/aa8955>

Harman, C. E., Schwieterman, E. W., Schottelkotte, J. C., & Kasting, J. F. (2015). Abiotic O<sub>2</sub> levels on planets around F, G, K, and M stars: possible false positives for life?. *The Astrophysical Journal*, 812(2), 137. <https://doi.org/10.1088/0004-637X/812/2/137>

Hatzes, A. P. (2010). A European Roadmap for exoplanets. *Proceedings of the International Astronomical Union*, 6(S276), 316-323. <https://doi.org/10.1017/S1743921311020382>

Hitchcock, D. R., & Lovelock, J. E. (1967). Life detection by atmospheric analysis. *Icarus*, 7(1-3), 149-159. [https://doi.org/10.1016/0019-1035\(67\)90059-0](https://doi.org/10.1016/0019-1035(67)90059-0)

Kopparapu, R., et al. (2017). Habitable moist atmospheres on terrestrial planets near the inner edge of the habitable zone around M dwarfs. *The Astrophysical Journal*, 845(1), 5. <https://doi.org/10.3847/1538-4357/aa7cf9>

Kouveliotou, C., et al. (2014). Enduring quests-daring visions (NASA astrophysics in the next three decades). *arXiv preprint arXiv:1401.3741*. <https://arxiv.org/abs/1401.3741>

Krissansen-Totton, J., Bergsman, D. S., & Catling, D. C. (2016). On detecting biospheres from chemical thermodynamic disequilibrium in planetary atmospheres. *Astrobiology*, 16(1), 39-67. <https://doi.org/10.1089/ast.2015.1327>

Lehmer, O. R., Catling, D. C., Parenteau, M. N., & Hoehler, T. M. (2018). The productivity of oxygenic photosynthesis around cool, M dwarf stars. *The Astrophysical Journal*, 859(2), 171. <https://doi.org/10.3847/1538-4357/aac104>

Lovelock, J. E. (1965). A physical basis for life detection experiments. *Nature*, 207(997), 568-570. <https://www.nature.com/articles/207568a0>

Loyd, R. P., et al. (2018). The MUSCLES Treasury Survey. V. FUV Flares on Active and Inactive M Dwarfs. *The Astrophysical Journal*, 867(1), 71. <https://doi.org/10.3847/1538-4357/aae2bd>

- Lincowski, A. P., et al. (2018). Evolved Climates and Observational Discriminants for the TRAPPIST-1 Planetary System. *The Astrophysical Journal*, 867(1), 76. <https://doi.org/10.3847/1538-4357/aae36a>
- Luger, R., & Barnes, R. (2015). Extreme water loss and abiotic O<sub>2</sub> buildup on planets throughout the habitable zones of M dwarfs. *Astrobiology*, 15(2), 119-143. <https://doi.org/10.1089/ast.2014.1231>
- Luger, R., Barnes, R., Lopez, E., Fortney, J., Jackson, B., & Meadows, V. (2015). Habitable evaporated cores: transforming mini-Neptunes into super-Earths in the habitable zones of M dwarfs. *Astrobiology*, 15(1), 57-88. <https://doi.org/10.1089/ast.2014.1215>
- Lunine, J. I., et al., (2008), Worlds beyond: A strategy for the detection and characterization of exoplanets executive summary of a report of the exoplanet task force astronomy and astrophysics advisory committee Washington, DC June 23, 2008. *Astrobiology*, 8, 875. <https://doi.org/10.1089/ast.2008.0276>
- MacGregor, M. A., Weinberger, A. J., Wilner, D. J., Kowalski, A. F., & Cranmer, S. R. (2018). Detection of a millimeter flare from Proxima Centauri. *The Astrophysical Journal Letters*, 855(1), L2. <https://doi.org/10.3847/2041-8213/aaad6b>
- Meadows, V.S., (2017). Reflections on O<sub>2</sub> as a biosignature in exoplanetary atmospheres. *Astrobiology*, 17(10), pp.1022-1052. <https://doi.org/10.1089/ast.2016.1578>
- Morbidelli, A., et al., (2000). Source regions and timescales for the delivery of water to the Earth. *Meteoritics & Planetary Science*, 35(6), 1309-1320. <https://doi.org/10.1111/j.1945-5100.2000.tb01518.x>
- National Academies of Sciences (NAS), Engineering, and Medicine. (2018). *Exoplanet Science Strategy*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/25187>
- Olson, S. L., Reinhard, C. T., & Lyons, T. W. (2016). Limited role for methane in the mid-Proterozoic greenhouse. *Proceedings of the National Academy of Sciences*, 113(41), 11447-11452. <https://doi.org/10.1073/pnas.1608549113>
- Owen, J. E., & Mohanty, S. (2016). Habitability of terrestrial-mass planets in the HZ of M Dwarfs—I. H/He-dominated atmospheres. *Monthly Notices of the Royal Astronomical Society*, 459(4), 4088-4108. <https://doi.org/10.1093/mnras/stw959>
- Quanz, S. P., Crossfield, I., Meyer, M. R., Schmalzl, E., & Held, J. (2015). Direct detection of exoplanets in the 3–10 μm range with E-ELT/METIS. *International Journal of Astrobiology*, 14(2), 279-289. <https://doi.org/10.1017/S1473550414000135>

Ramirez, R. M., & Kaltenegger, L. (2014). The habitable zones of pre-main-sequence stars. *The Astrophysical Journal Letters*, 797(2), L25. <https://doi.org/10.1088/2041-8205/797/2/L25>

Ramirez, R. M., & Kaltenegger, L. (2018). A methane extension to the classical habitable zone. *The Astrophysical Journal*, 858(2), 72. <https://doi.org/10.3847/1538-4357/aab8fa>

Reinhard, C. T., Olson, S. L., Schwieterman, E. W., & Lyons, T. W. (2017). False negatives for remote life detection on ocean-bearing planets: lessons from the early Earth. *Astrobiology*, 17(4), 287-297. <https://doi.org/10.1089/ast.2016.1598>

Richey-Yowell, T. et al., (2019). HAZMAT. V. The Ultraviolet and X-ray Evolution of K Stars. *The Astrophysical Journal* 872, 17 <https://doi.org/10.3847/1538-4357/aafa74>

Rogers, L. A. (2015). Most 1.6 Earth-radius planets are not rocky. *The Astrophysical Journal*, 801(1), 41. <https://doi.org/10.1088/0004-637X/801/1/41>

Seager, S., et al. (2015). The Exo-S probe class starshade mission. In *Techniques and Instrumentation for Detection of Exoplanets VII* (Vol. 9605, p. 96050W). International Society for Optics and Photonics. <https://doi.org/10.1117/12.2190378>

Shkolnik, E. L., & Barman, T. S. (2014). HAZMAT. I. The evolution of far-UV and near-UV emission from early M stars. *The Astronomical Journal*, 148(4), 64. <https://doi.org/10.1088/0004-6256/148/4/64>

Schneider, A. C., & Shkolnik, E. L. (2018). HAZMAT. III. The UV Evolution of Mid-to Late-M Stars with GALEX. *The Astronomical Journal*, 155(3), 122. <https://doi.org/10.3847/1538-3881/aaaa24>

Schwieterman, E. W., et al. (2018). Exoplanet biosignatures: a review of remotely detectable signs of life. *Astrobiology*, 18(6), 663-708. <https://doi.org/10.1089/ast.2017.1729>

Schwieterman, E. W., et al. (2019). A Limited Habitable Zone for Complex Life. <https://arxiv.org/abs/1902.04720>

Stapelfeldt, et al.(2014). Exo-C: a probe-scale space mission to directly image and spectroscopically characterize exoplanetary systems using an internal coronagraph. In *Space Telescopes and Instrumentation 2014: Optical, Infrared, and Millimeter Wave* (Vol. 9143, p. 91432K). International Society for Optics and Photonics. <https://doi.org/10.1117/12.2057115>

Tian, F., France, K., Linsky, J. L., Mauas, P. J., & Vieytes, M. C. (2014). High stellar FUV/NUV ratio and oxygen contents in the atmospheres of potentially habitable planets. *Earth and Planetary Science Letters*, 385, 22-27. <https://doi.org/10.1016/j.epsl.2013.10.024>

Tian, F. and Ida, S., 2015. Water contents of Earth-mass planets around M dwarfs. *Nature Geoscience*, 8(3), p.177. <https://www.nature.com/articles/ngeo2372>

Wang, Z., & Becker, H. (2013). Ratios of S, Se and Te in the silicate Earth require a volatile-rich late veneer. *Nature*, 499(7458), 328. <https://www.nature.com/articles/nature12285>

West, A. A., et al.(2015). An Activity–Rotation Relationship and Kinematic Analysis of Nearby Mid-to-Late-Type M Dwarfs. *The Astrophysical Journal*, 812(1), 3. <https://doi.org/10.1088/0004-637X/812/1/3>

Wordsworth, R., & Pierrehumbert, R. (2014). Abiotic oxygen-dominated atmospheres on terrestrial habitable zone planets. *The Astrophysical Journal Letters*, 785(2), L20. <https://doi.org/10.1088/2041-8205/785/2/L20>

Yang, J., Cowan, N. B., & Abbot, D. S. (2013). Stabilizing cloud feedback dramatically expands the habitable zone of tidally locked planets. *The Astrophysical Journal Letters*, 771(2), L45. <https://doi.org/10.1088/2041-8205/771/2/L45>