

Astro2020 Science White Paper

Radio Pulsar Populations

- Thematic Areas:**
- Planetary Systems
 - Star and Planet Formation
 - Formation and Evolution of Compact Objects
 - Cosmology and Fundamental Physics
 - Stars and Stellar Evolution
 - Resolved Stellar Populations and their Environments
 - Galaxy Evolution
 - Multi-Messenger Astronomy and Astrophysics

Principal Author:

Name: Duncan Lorimer
Institution: West Virginia University
Email: duncan.lorimer@mail.wvu.edu
Phone: 304-290-0417

Co-authors:

Nihan Pol (West Virginia University)
Kaustubh Rajwade (University of Manchester)
Kshitij Aggarwal (West Virginia University)
Devansh Agarwal (West Virginia University)
Jay Strader (Michigan State University)
Natalia Lewandowska (West Virginia University)
David Kaplan (University of Wisconsin Milwaukee)
Tyler Cohen (New Mexico Tech)
Paul Demorest (National Radio Astronomy Observatory)
Emmanuel Fonseca (McGill University)
Shami Chatterjee (Cornell University)

Abstract: Our understanding of the neutron star population is informed to a great degree by large-scale surveys that have been carried out by radio facilities during the past fifty years. We summarize some of the recent breakthroughs in our understanding of the radio pulsar population, and look ahead to future yields from upcoming experiments planned for the next decade. By the end of the 2020s, we anticipate a much more complete census of the Galactic population and being able to probe the populations of radio-emitting neutron stars more effectively in external galaxies. Among the anticipated discoveries are pulsar–black hole binary systems that will provide further tests of strong-field gravity, as well as large numbers of millisecond pulsars that are crucial to enhancing the sensitivity of timing arrays for low-frequency gravitational waves.

1 Why Study Pulsar Populations?

Radio pulsars, rapidly spinning highly magnetized neutron stars, have long been known as excellent tools to study fundamental physics and astronomy [1, 2]. Of particular interest are “millisecond pulsars” (MSPs) which rotate with frequencies of several hundred Hz with atomic clock-like precision. MSPs are frequently used as tools to investigate a variety of topics, including: the physics of matter at supranuclear densities [3, 4], detection of low-frequency gravitational waves [5], probes of general relativity [6], interstellar weather [7], globular cluster astrophysics [8], extrasolar planets [9, 10] and planetary physics [11]. Three of the many key astrophysical questions to be tackled in the upcoming decade through large-scale surveys and detailed modeling are outlined below.

1. How fast can a neutron star spin? On theoretical grounds, the maximum spin rate is governed by the density profile of neutron stars [12]. Softer profiles allow for the existence of more rapidly spinning neutron stars, and observations of any such objects would rule out the stiffer profiles. X-ray observations [13] suggest that the maximum spin frequency may be close to 700 Hz. Observationally, the fastest spinning neutron star is the 716 Hz pulsar in Terzan 5 [14]. Do neutron stars exist with > 1 kHz spin rates? Future surveys could definitively answer this question.

2. How many pulsars are there? Studies of the spatial distribution of normal pulsars have shown that pulsars generally trace the radial distribution of massive stars in our Galaxy, but have a scale height above the plane of several hundred parsecs [15]. Although the vertical distribution of MSPs has been studied locally [16], the radial density of MSPs is poorly understood. This topic is of great interest to: (i) our ongoing efforts [17] to optimize future surveys to find MSPs for gravitational wave detection and characterization by pulsar timing arrays; (ii) understanding the MSP contribution to the diffuse Galactic gamma-ray emission [18].

3. How are pulsars formed? Pulsars are superb probes of binary star evolution. The existence of isolated MSPs and unusual binary/planetary systems implies that other progenitors are required to explain the MSP population we see beyond those predicted by the standard model. Of key importance is determining the birth rates of the various sub-samples of MSPs and comparing them to alternative formation routes (e.g. the accretion-induced collapse of a white dwarf [19]). Many of the details of the assumptions (e.g. mass transfer physics and neutron star birth velocities) are still not well understood. Improved knowledge of these areas would help studies of source classes for gravitational wave emission from compact binary inspirals [20].

Despite over 50 years of study, it is not well known how pulsars radiate. There is a wide variety of emission behavior from pulse to pulse through nulling, mode-changing, giant pulses and sub-pulse drifting phenomena [21, 22, 23, 24, 25, 26]. Closely linked phenomena are the rotating radio transients [27] and the intermittent pulsars [28, 29, 30]. While the rotating radio transients emit only occasional individual pulses spaced anywhere between seconds to years, intermittent pulsars switch on and remain steady before switching off again over timescales of days to years. Over these longer timescales, where it is possible to measure the spin period derivative precisely, it is observed that the “on” and “off” states for the intermittent pulsars are associated with two different spin-down rates! A similar behavior was seen in 17 non-intermittent pulsars [31] which switch between different spin-down states in an apparently quasi-periodic fashion, sometimes accompanied by correlated pulse profile changes. The mixture of spin-down states could even account for the timing noise variations are seen in many pulsars and suggest that most, if not all, normal pulsars have multiple spin-down states indicating global changes in magnetospheric current densities.

2 Modeling Pulsar Populations

Inferring the properties of the underlying, as opposed to the observed, pulsar population is a difficult problem. Not only does the detection process have to be accurately accounted for to account for observational biases, the non-homogenous properties of the interstellar medium need to be taken into account. Much of the progress has come from detailed Monte Carlo simulations of pulsar populations that create artificial pulsars that satisfy the criteria for detection, it is possible to generate and optimize models for the pulsar population which inform us about the underlying distribution functions and make predictions for future survey yields. Generally, the Monte Carlo simulations follow two basic strategies. In the “snapshot” approach, no assumptions are made concerning the prior evolution of pulsars. Instead, populations are generated according to various distribution functions (typically in Galactocentric radius, R , height with respect to the plane, z , spin period, P and luminosity, L) which are optimized to find the best match to the sample.




Figure 1: P - \dot{P} diagram showing Monte Carlo pulsars detected in models of surveys carried out with the Parkes telescope (bold points). P - \dot{P} values that were taken from the pulsar catalog (smaller points). Histograms of the P and \dot{P} distributions are also shown, with grey bars for catalog sources, and with solid lines for the simulated pulsars. Figure is taken from Bates et al. [32].

Alternatively, as shown in Fig. 1, one may carry out “evolution” approaches where the model pulsars are evolved forward in time from a set of initial distributions. Software to carry out both approaches has been developed by a number of groups and some of this is freely available. For example, the `PSRPOP` software package [32] has modules to carry out both the snapshot and evolution approaches. Although many earlier studies did not attempt to rigorously optimize model parameters [33], in recent years these approaches have become more and more exhaustive and employed Markov-Chain Monte Carlo methods to estimate the density of pulsars in phase space and optimize these to find marginalized probability density functions for various model parameters [34, 35] and to compute Bayes factors to compare different families of models.

3 Pulsar Demography: Recent Advances

Inspired by earlier studies [36, 37], the snapshot approach was applied to the normal pulsar population [15] to derive best-fitting probability density functions in R , L , z and P for the present-day population of objects. One result of this work was that the radial distribution of pulsars could not be decoupled from the radial distribution of free electrons in the pulsar distribution. For the evolution approach on the normal population, a highly cited study is the work of Faucher-Giguère & Kaspi [33] who generated excellent fits to the pulsar $P-\dot{P}$ diagram (Fig. 1) using a model in which the luminosity has a power-law dependence on P and \dot{P} . In addition, the luminosity function of the present-day pulsar population appears to be log-normal in form. The snapshot technique can be modified [38] such that it can be combined with the census of observed neutron star-white dwarf and double neutron star binary systems to calculate their respective merger detection rates for ground and space-based gravitational wave detectors [39, 40, 41].

Recent progress has been made in our understanding of the evolution of the magnetic inclination angle, α , for normal pulsars. Following earlier work [42], more detailed modeling following the evolutionary framework outlined above by Johnston & Karastergiou [43] demonstrated that an exponential decay of α on a 10^7 yr time scale can provide an excellent match to the $P-\dot{P}$ diagram for a pulsar birth rate of one per century and an initial spin period of 20 ms. Historians of pulsar statistics will be aware that the debate surrounding initial spins of normal pulsars, their magnetic field and luminosity evolution, as well as the role of α in the spin-down evolution model, has been a contentious issue. The new developments from this study, however, provide a framework for more accurately modeling the evolution of the pulsar spin-down than has been previously possible. Theoretical progress in the past few years has arisen from detailed magnetohydrodynamical modeling [44]. As a result, most of the pulsar evolution codes are now moving away from the simple magnetic dipole model [45] that has been used almost exclusively up until around 2010.

Our understanding of the Galactic population of MSPs and, more generally, recycled pulsars over the past decade has been significantly improved. This is thanks, in large part, to a great increase in sample size provided by a number of large-scale surveys of the Galactic disk at Parkes [46], Green Bank [47], Arecibo [48], Effelsberg [49] and the Low-Frequency Array [50]. These surveys have greatly increased the sample size such that the number of Galactic MSPs now outnumber their counterparts in globular clusters. A recent study by Gonthier et al. [35] combined radio surveys with results from the Fermi point source catalog [51] to provide a comprehensive model of the radio and gamma-ray MSP population.

Strong evidence for selection bias in MSPs comes from γ -ray observations with *Fermi*. Among the most exciting discoveries of *Fermi* has been that non-accreting pulsars are nearly ubiquitous emitters of GeV γ -rays [52]. In addition to the physical insight possible by linking radio and high-energy observations of individual systems, radio timing observations of previously unassociated *Fermi* sources have allowed the discovery of many new pulsars. In particular, there has been an enormous increase in the number of binary millisecond pulsars that ablate their hydrogen-rich companions. These are typically classified as either “black widow” or “redback” systems depending on whether the companion is less or more massive than $\sim 0.1M_{\odot}$ [53]. The pulsed radio emission from these systems is sometimes eclipsed due to ionized material associated with the companion, but this material apparently does not affect the detectability of the GeV emission. Thus, while existing pulsar catalogs are biased against systems with hydrogen-rich companions, follow-up of unassociated *Fermi* sources has proven an effective way to discover and characterize these binaries

[54]. These “spider” binaries are important to improve our understanding of the process by which pulsars are recycled, and to determine long-sought values such as the maximum mass of a neutron star. It is now clear that these binaries host more massive neutron stars than the canonical $1.4M_{\odot}$, with median masses of $1.8M_{\odot}$ [55], and several systems having minimum masses of $> 1.9M_{\odot}$. The discovery and follow-up of more such binaries, which is best accomplished through optical, X-ray, γ -ray, and radio observations, offers one of the best paths to confirm neutron stars with masses $> 2M_{\odot}$ to determine the maximum neutron star mass.

Two further outstanding questions are the population of pulsars in the inner Galaxy and Galactic halo. In the central regions, there is intense interest in constraining this population both from the point of view of finding pulsars in orbit around SgrA* [56] and from investigating whether the Galactic Center gamma-ray excess is due to an ensemble of millisecond pulsars [57]. Due to the impacts of pulse dispersion and scattering, as well as the significant inverse square law distance penalty to the Galactic Center, searches have so far had limited sensitivity. A recent study taking these factors into account [58] suggests that the optimal frequency for future surveys of this region is 9–13 GHz. In the Galactic halo, a recent analysis shows that the fraction of the population bound therein depends critically on the magnetic angle evolution and birth velocity distribution [59].

The pulsar content in Galactic globular clusters has been studied extensively over the past thirty years. Recent studies have made use of constraints from the luminosity function inferred from the Galactic disk population to make predictions on the various types of pulsars. Highlights from this work include constraints on the highly anomalous population of young (non-recycled) pulsars in globular clusters [60] as well a census of the entire population [61]. The key relationship between overall pulsar abundance appears to be the two-body encounter rate [61, 62]. This leads to a number of predictions for cluster searches with new facilities now coming online.

Currently, the most distant pulsars known are in the Large and Small Magellanic Clouds [63, 64]. The relatively small sample size at present has hampered attempts to model the pulsar content in these galaxies and has so far prohibited detailed comparisons with the Milky Way and globular cluster population. The next decade is likely to result in the discovery of a substantial number of extragalactic pulsars, which can be used to probe the Galactic halo and circumgalactic medium.

4 Key Advances: Upcoming Surveys

Many facilities are set to transform our understanding of the radio pulsar population over the next decade. Most notable are the Five Hundred Meter Aperture Radio Telescope (FAST), the Canadian H I Intensity Mapping Experiment (CHIME), Hydrogen Intensity and Real-time Analysis eXperiment (HIRAX), MeerKAT (Karoo Array Telescope), the Deep Synoptic Array (DSA), the Australian Square Kilometer Array Pathfinder (ASKAP), the next generation Very Large Array (ngVLA) and the Square Kilometer Array (SKA). Existing facilities such as Arecibo (currently being outfitted with a 40 beam L-band system), Parkes, Effelsberg, the upgraded Westerbork Synthesis Radio Telescope array (WSRT), the Green Bank Telescope (GBT), the VLA, the upgraded Giant Metre Wave Radio Telescope (uGMRT), the Low Frequency Array (LOFAR), and the Long Wavelength Array (LWA) will also be very important.

All these facilities will be carrying out pulsar surveys which are in varying states of plan-

Telescope	Frequency band (GHz)	Gain (K/Jy)	Integration (s)	Sky coverage (°)	N_{normal}	N_{MSP}
Arecibo	1.3–1.8	10	600	$0 < \delta < +38$	4200	850
CHIME	0.4–0.8	2	900	$\delta > -20$	5600	650
MeerKAT	1.1–1.8	2.6	300	$-90 < \delta < +40$	12000	1600
FAST	1.1–1.9	20	20	$-20 < \delta < +60$	3400	1800
ngVLA	1.2–3.5	7.6	240	$-15 < l < 265; b < 1$	2200	100

Table 1: Summary of `PsRPopPy` runs showing the predicted sample sizes of both normal pulsars (N_{normal}) and MSPs (N_{MSP}) obtained from the online simulator for putative surveys planned for the coming decade. We do not account for overlapping detections, or currently known pulsars.

ning and scope. We provide a brief overview of the expected results¹ for some of them based on `PsRPopPy`. Survey strategies vary greatly, depending on the characteristics of each instrument. In Table 1, we summarize the number of normal and MSPs detectable by putative searches with each of the above facilities. The numbers are illustrative in the sense that they are based on the entire sky accessible from each telescope and integration times that may take several years to cover in some cases. In addition, they do not account for the number of pulsars currently known². However, they do provide a measure of each telescope’s potential as a probe of the Galactic pulsar population and can be scaled and refined accordingly in the future as the experiments are being carried out. As these estimates demonstrate, the radio pulsar population will increase substantially in the coming decade. We anticipate a Galactic sample size approaching 10^4 pulsars and $> 10^3$ MSPs. Excellent prospects for finding Galactic Center pulsars come from the ngVLA and SKA [66].

5 Concluding Remarks

The past 50 years of pulsar astrophysics has informed us that we are only scratching the surface of the neutron star zoo [67]. Some of the landmark discoveries during this time have included the original double neutron star binary [68], millisecond pulsars [69], pulsar planetary systems [9], the double pulsar [70, 71], radio magnetars [72], fast radio bursts [73], ultra-light degenerate companions [74, 75], and triple systems [76, 77, 78, 79, 80]. With such a large yield expected by the upcoming surveys, further rare and exotic discoveries are anticipated. High among these would be the discovery of pulsar–black hole binary systems. These are anticipated in the Galactic field [81], but also around Sgr A* [82] and in globular cluster systems [83]. Such systems could provide highly sensitive test beds of general relativity that go well beyond the already impressive results [84]. The large numbers of MSPs anticipated will be crucial to the sensitivities of the pulsar timing arrays as they move into an era of low-frequency gravitational wave detection [85].

¹Yields for any survey can be estimated using the `PsRPopPy` simulator at <http://psrpop.phys.wvu.edu>.

²As of March 2019, 2659 radio pulsars are currently listed in the Australia Telescope National Facility (ATNF) catalog [65] which can be accessed at <http://www.atnf.csiro.au/research/pulsar/psrcat>.

References

- [1] R. D. Blandford, A. Hewish, A. G. Lyne, and L. Mestel. Pulsars as Physical Laboratories. **341** 1, 1992.
- [2] D. R. Lorimer and M. Kramer. *Handbook of Pulsar Astronomy*. Cambridge University Press, 2005.
- [3] P. B. Demorest, T. Pennucci, S. M. Ransom, M. S. E. Roberts, and J. W. T. Hessels. A two-solar-mass neutron star measured using Shapiro delay. **467** 1081, 2010. doi:10.1038/nature09466. 1010.5788.
- [4] F. Özel, D. Psaltis, S. Ransom, P. Demorest, and M. Alford. The Massive Pulsar PSR J1614-2230: Linking Quantum Chromodynamics, Gamma-ray Bursts, and Gravitational Wave Astronomy. **724** L199, 2010. doi:10.1088/2041-8205/724/2/L199. 1010.5790.
- [5] F. A. Jenet, G. B. Hobbs, W. van Straten, R. N. Manchester, M. Bailes, J. P. W. Verbiest, R. T. Edwards, A. W. Hotan, J. M. Sarkissian, and S. M. Ord. Upper Bounds on the Low-Frequency Stochastic Gravitational Wave Background from Pulsar Timing Observations: Current Limits and Future Prospects. **653** 1571, 2006. doi:10.1086/508702. astro-ph/0609013.
- [6] M. Kramer, A. G. Lyne, M. Burgay, A. Possenti, R. N. Manchester, F. Camilo, M. A. McLaughlin, D. R. Lorimer, N. D’Amico, B. C. Joshi, J. Reynolds, and P. C. C. Freire. The double pulsar - a new testbed for relativistic gravity. pages 59–65. Astro-ph/0405179.
- [7] R. Ramachandran, P. Demorest, D. C. Backer, I. Cognard, and A. Lommen. Interstellar Plasma Weather Effects in Long-Term Multifrequency Timing of Pulsar B1937+21. **645** 303, 2006. doi:10.1086/500634. arXiv:astro-ph/0601242.
- [8] F. Camilo and F. A. Rasio. Pulsars in globular clusters. page 147.
- [9] A. Wolszczan and D. A. Frail. A planetary system around the millisecond pulsar PSR 1257+12. **355** 145, 1992.
- [10] M. Bailes, S. D. Bates, V. Bhlerao, N. D. R. Bhat, M. Burgay, S. Burke-Spolaor, N. D’Amico, S. Johnston, M. J. Keith, M. Kramer, S. R. Kulkarni, L. Levin, A. G. Lyne, S. Milia, A. Possenti, L. Spitler, B. Stappers, and W. van Straten. Transformation of a Star into a Planet in a Millisecond Pulsar Binary. **333** 1717, 2011. doi:10.1126/science.1208890. 1108.5201.
- [11] D. J. Champion, G. B. Hobbs, R. N. Manchester, R. T. Edwards, D. C. Backer, M. Bailes, N. D. R. Bhat, S. Burke-Spolaor, W. Coles, P. B. Demorest, R. D. Ferdman, W. M. Folkner, A. W. Hotan, M. Kramer, A. N. Lommen, D. J. Nice, M. B. Purver, J. M. Sarkissian, I. H. Stairs, W. van Straten, J. P. W. Verbiest, and D. R. B. Yardley. Measuring the Mass of Solar System Planets Using Pulsar Timing. **720** L201, 2010. doi:10.1088/2041-8205/720/2/L201. 1008.3607.
- [12] J. H. Lattimer and M. Prakash. The physics of neutron stars. **304** 536, 2004.

- [13] D. Chakrabarty, E. H. Morgan, M. P. Muno, D. K. Galloway, R. Wijnands, M. van der Klis, and C. B. Markwardt. Nuclear-powered millisecond pulsars and the maximum spin frequency of neutron stars. **424** 42, 2003.
- [14] J. Hessels, S. Ransom, I. Stairs, P. Freire, V. Kaspi, and F. Camilo. A radio pulsar spinning at 716 hz. *Science*, **311** 1901, 2006. doi:10.1126/science.1123430. arXiv:astro-ph/0601337.
- [15] D. R. Lorimer, A. J. Faulkner, A. G. Lyne, R. N. Manchester, M. Kramer, M. A. McLaughlin, G. Hobbs, A. Possenti, I. H. Stairs, F. Camilo, M. Burgay, N. D’Amico, A. Corongiu, and F. Crawford. The Parkes Multibeam Pulsar Survey - VI. Discovery and timing of 142 pulsars and a Galactic population analysis. **372** 777, 2006. arXiv:astro-ph/0607640.
- [16] D. R. Lorimer. Pulsar Statistics – II. The local low-mass binary pulsar population. **274** 300, 1995.
- [17] M. McLaughlin. The North American Nanohertz Observatory for Gravitational Waves. *Class. Quantum Grav.*, page 224008, 2013.
- [18] C.-A. Faucher-Giguère and A. Loeb. The pulsar contribution to the gamma-ray background. *Journal of Cosmology and Astroparticle Physics*, **1** 5, 2010. doi:10.1088/1475-7516/2010/01/005. 0904.3102.
- [19] R. Narayan and R. Popham. Angular momentum of accreting white dwarfs: Implications for millisecond pulsar formation. **346** L25, 1989.
- [20] R. O’Shaughnessy, C. Kim, T. Fragos, V. Kalogera, and K. Belczynski. Constraining Population Synthesis Models via the Binary Neutron Star Population. **633** 1076, 2005. doi:10.1086/468180. arXiv:astro-ph/0504479.
- [21] D. C. Backer. Pulsar nulling phenomena. **228** 42, 1970.
- [22] D. C. Backer. Peculiar pulse burst in PSR 1237+25. **228** 1297, 1970.
- [23] D. C. Backer. Correlation of subpulse structure in a sequence of pulses from pulsar psr 1919+21. **227** 692, 1970.
- [24] D. C. Backer. Correlation of subpulse structure in psr 1237+25. **228** 752, 1970.
- [25] V. A. Soglasnov, M. V. Popov, N. Bartel, W. Cannon, A. Y. Novikov, V. I. Kondratiev, and V. I. Altunin. Giant Pulses from PSR B1937+21 with Widths =15 Nanoseconds and $T_b=5\times 10^{39}$ K, the Highest Brightness Temperature Observed in the Universe. , **616** 439, 2004. doi:10.1086/424908. astro-ph/0408285.
- [26] D. H. Staelin and E. C. Reifenstein, III. Pulsating Radio Sources near the Crab Nebula. *Science*, **162** 1481, 1968. doi:10.1126/science.162.3861.1481.
- [27] M. A. McLaughlin, A. G. Lyne, D. R. Lorimer, M. Kramer, A. J. Faulkner, R. N. Manchester, J. M. Cordes, A. Possenti, F. Camilo, G. Hobbs, I. H. Stairs, N. D’Amico, and J. T. O’Brien. Transient radio bursts from rotating neutron stars. **439** 817, 2006.

- [28] M. Kramer, A. G. Lyne, J. T. O'Brien, C. A. Jordan, and D. R. Lorimer. A Periodically Active Pulsar Giving Insight into Magnetospheric Physics. *Science*, **312** 549, 2006. arXiv:astro-ph/0604605.
- [29] F. Camilo, S. M. Ransom, S. Chatterjee, S. Johnston, and P. Demorest. PSR J1841-0500: A Radio Pulsar That Mostly is Not There. **746** 63, 2012. doi:10.1088/0004-637X/746/1/63.1111.5870.
- [30] D. R. Lorimer, A. G. Lyne, M. A. McLaughlin, M. Kramer, G. G. Pavlov, and C. Chang. Radio and X-Ray Observations of the Intermittent Pulsar J1832+0029. **758** 141, 2012. doi:10.1088/0004-637X/758/2/141.1208.6576.
- [31] A. Lyne, G. Hobbs, M. Kramer, I. Stairs, and B. Stappers. Switched Magnetospheric Regulation of Pulsar Spin-Down. *Science*, **329** 408, 2010. doi:10.1126/science.1186683.1006.5184.
- [32] S. D. Bates, D. R. Lorimer, A. Rane, and J. Swiggum. PsrPopPy: An open-source package for pulsar population simulations. 2013. Submitted (arXiv:1311.3247), 1311.3247.
- [33] C.-A. Faucher-Giguère and V. M. Kaspi. Birth and Evolution of Isolated Radio Pulsars. **643** 332, 2006. doi:10.1086/501516. astro-ph/0512585.
- [34] M. Cieřlar, T. Bulik, and S. Osłowski. Markov chain Monte Carlo population synthesis of single radio pulsars in the Galaxy. arXiv e-prints, 2018. 1803.02397.
- [35] P. L. Gonthier, A. K. Harding, E. C. Ferrara, S. E. Frederick, V. E. Mohr, and Y.-M. Koh. Population Syntheses of Millisecond Pulsars from the Galactic Disk and Bulge. , **863** 199, 2018. doi:10.3847/1538-4357/aad08d.1806.11215.
- [36] J. H. Taylor and R. N. Manchester. Galactic distribution and evolution of pulsars. **215** 885, 1977.
- [37] A. G. Lyne, R. N. Manchester, and J. H. Taylor. The Galactic population of pulsars. **213** 613, 1985.
- [38] C. Kim, V. Kalogera, and D. R. Lorimer. The probability distribution of binary pulsar coalescence rates. i. double neutron star systems in the galactic field. **584** 985, 2003.
- [39] C. Kim, V. Kalogera, D. R. Lorimer, and T. White. The Probability Distribution Of Binary Pulsar Coalescence Rates. II. Neutron Star-White Dwarf Binaries. , **616** 1109, 2004. doi:10.1086/424954. astro-ph/0402162.
- [40] C. Kim, B. B. P. Perera, and M. A. McLaughlin. Implications of PSR J0737-3039B for the Galactic NS-NS binary merger rate. , **448** 928, 2015. doi:10.1093/mnras/stu2729.1308.4676.
- [41] N. Pol, M. McLaughlin, and D. R. Lorimer. Future Prospects for Ground-based Gravitational-wave Detectors: The Galactic Double Neutron Star Merger Rate Revisited. , **870** 71, 2019. doi:10.3847/1538-4357/aaf006.1811.04086.

- [42] P. Weltevrede and S. Johnston. The population of pulsars with interpulses and the implications for beam evolution. , **387** 1755, 2008. doi:10.1111/j.1365-2966.2008.13382.x. 0804.4318.
- [43] S. Johnston and A. Karastergiou. Pulsar braking and the \dot{P} diagram. , **467** 3493, 2017. doi:10.1093/mnras/stx377. 1702.03616.
- [44] A. Spitkovsky. Time-dependent Force-free Pulsar Magnetospheres: Axisymmetric and Oblique Rotators. , **648** L51, 2006. doi:10.1086/507518. astro-ph/0603147.
- [45] J. E. Gunn and J. P. Ostriker. On the nature of pulsars. III. Analysis of observations. **160** 979, 1970.
- [46] R. N. Manchester, A. G. Lyne, F. Camilo, J. F. Bell, V. M. Kaspi, N. D’Amico, N. P. F. McKay, F. Crawford, I. H. Stairs, A. Possenti, D. J. Morris, and D. C. Sheppard. The Parkes multi-beam pulsar survey - i. observing and data analysis systems, discovery and timing of 100 pulsars. **328** 17, 2001.
- [47] R. S. Lynch. The Green Bank North Celestial Cap Pulsar Survey: Status and Future. In P. Weltevrede, B. B. P. Perera, L. L. Preston, and S. Sanidas, editors, *Pulsar Astrophysics the Next Fifty Years*, volume 337 of *IAU Symposium*, pages 13–16. 2018. doi: 10.1017/S174392131700953X.
- [48] J. M. Cordes, P. C. C. Freire, D. R. Lorimer, F. Camilo, D. J. Champion, D. J. Nice, R. Ramachandran, J. W. T. Hessels, W. Vlemmings, J. van Leeuwen, S. M. Ransom, N. D. R. Bhat, Z. Arzoumanian, M. A. McLaughlin, V. M. Kaspi, L. Kasian, J. S. Deneva, B. Reid, S. Chatterjee, J. L. Han, D. C. Backer, I. H. Stairs, A. A. Deshpande, and C.-A. Faucher-Giguère. Arecibo Pulsar Survey Using ALFA. I. Survey Strategy and First Discoveries. **637** 446, 2006. astro-ph/0509732.
- [49] E. Barr. The Northern-sky High Time Resolution Universe Pulsar Survey. In M. Burgay, N. D’Amico, P. Esposito, A. Pellizzoni, and A. Possenti, editors, *American Institute of Physics Conference Series*, volume 1357 of *American Institute of Physics Conference Series*, pages 52–53. 2011. doi:10.1063/1.3615075.
- [50] T. Coenen, J. van Leeuwen, J. W. T. Hessels, B. W. Stappers, V. I. Kondratiev, A. Alexov, R. P. Breton, A. Bilous, S. Cooper, H. Falcke, R. A. Fallows, V. Gajjar, J.-M. Grießmeier, T. E. Hassall, A. Karastergiou, E. F. Keane, M. Kramer, M. Kuniyoshi, A. Noutsos, S. Osłowski, M. Pilia, M. Serylak, C. Schrijvers, C. Sobey, S. ter Veen, J. Verbiest, P. Weltevrede, S. Wijnholds, K. Zagkouris, A. S. van Amesfoort, J. Anderson, A. Asgekar, I. M. Avruch, M. E. Bell, M. J. Bentum, G. Bernardi, P. Best, A. Bonafede, F. Breitling, J. Broderick, M. Brüggen, H. R. Butcher, B. Ciardi, A. Corstanje, A. Deller, S. Duscha, J. Eislöffel, R. Fender, C. Ferrari, W. Frieswijk, M. A. Garrett, F. de Gasperin, E. de Geus, A. W. Gunst, J. P. Hamaker, G. Heald, M. Hoft, A. van der Horst, E. Juette, G. Kuper, C. Law, G. Mann, R. McFadden, D. McKay-Bukowski, J. P. McKean, H. Munk, E. Orru, H. Paas, M. Pandey-Pommier, A. G. Polatidis, W. Reich, A. Renting, H. Röttgering, A. Rowlinson, A. M. M. Scaife, D. Schwarz, J. Sluman, O. Smirnov, J. Swinbank, M. Tagger, Y. Tang, C. Tasse, S. Thoudam, C. Toribio,

- R. Vermeulen, C. Vocks, R. J. van Weeren, O. Wucknitz, P. Zarka, and A. Zensus. The LO-FAR pilot surveys for pulsars and fast radio transients. , **570** A60, 2014. doi:10.1051/0004-6361/201424495. 1408.0411.
- [51] A. A. Abdo, M. Ackermann, M. Ajello, A. Allafort, E. Antolini, W. B. Atwood, M. Axelsson, L. Baldini, J. Ballet, G. Barbiellini, and et al. Fermi Large Area Telescope First Source Catalog. , **188** 405, 2010. doi:10.1088/0067-0049/188/2/405. 1002.2280.
- [52] A. A. Abdo, M. Ajello, A. Allafort, L. Baldini, J. Ballet, G. Barbiellini, M. G. Baring, D. Bastieri, A. Belfiore, R. Bellazzini, and et al. The Second Fermi Large Area Telescope Catalog of Gamma-Ray Pulsars. , **208** 17, 2013. doi:10.1088/0067-0049/208/2/17. 1305.4385.
- [53] M. S. E. Roberts. Surrounded by spiders! New black widows and redbacks in the Galactic field. In J. van Leeuwen, editor, *Neutron Stars and Pulsars: Challenges and Opportunities after 80 years*, volume 291 of *IAU Symposium*, pages 127–132. 2013. doi:10.1017/S174392131202337X. 1210.6903.
- [54] P. S. Ray, A. A. Abdo, D. Parent, D. Bhattacharya, B. Bhattacharyya, F. Camilo, I. Cognard, G. Theureau, E. C. Ferrara, A. K. Harding, D. J. Thompson, P. C. C. Freire, L. Guillemot, Y. Gupta, J. Roy, J. W. T. Hessels, S. Johnston, M. Keith, R. Shannon, M. Kerr, P. F. Michelson, R. W. Romani, M. Kramer, M. A. McLaughlin, S. M. Ransom, M. S. E. Roberts, P. M. Saz Parkinson, M. Ziegler, D. A. Smith, B. W. Stappers, P. Weltevrede, and K. S. Wood. Radio Searches of Fermi LAT Sources and Blind Search Pulsars: The Fermi Pulsar Search Consortium. arXiv e-prints, arXiv:1205.3089, 2012. 1205.3089.
- [55] J. Strader, S. Swihart, L. Chomiuk, A. Bahramian, C. Britt, C. C. Cheung, K. Dage, J. Halpern, K.-L. Li, R. P. Mignani, J. A. Orosz, M. Peacock, R. Salinas, L. Shishkovsky, and E. Tremou. Optical Spectroscopy and Demographics of Redback Millisecond Pulsar Binaries. , **872** 42, 2019. doi:10.3847/1538-4357/aafbaa. 1812.04626.
- [56] D. Psaltis, N. Wex, and M. Kramer. A Quantitative Test of the No-hair Theorem with Sgr A* Using Stars, Pulsars, and the Event Horizon Telescope. , **818** 121, 2016. doi:10.3847/0004-637X/818/2/121. 1510.00394.
- [57] G. Fragione, V. Pavlík, and S. Banerjee. Neutron stars and millisecond pulsars in star clusters: implications for the diffuse gamma-radiation from the Galactic Centre. , **480** 4955, 2018. doi:10.1093/mnras/sty2234. 1804.04856.
- [58] K. M. Rajwade, D. R. Lorimer, and L. D. Anderson. Detecting pulsars in the Galactic Centre. , **471** 730, 2017. doi:10.1093/mnras/stx1661. 1611.06977.
- [59] K. Rajwade, J. Chennamangalam, D. Lorimer, and A. Karastergiou. The Galactic halo pulsar population. , **479** 3094, 2018. doi:10.1093/mnras/sty1695. 1802.04690.
- [60] J. Boyles, D. R. Lorimer, P. J. Turk, R. Mnatsakanov, R. S. Lynch, S. M. Ransom, P. C. Freire, and K. Belczynski. Young Radio Pulsars in Galactic Globular Clusters. **742** 51, 2011.

- [61] P. J. Turk and D. R. Lorimer. An empirical Bayesian analysis applied to the globular cluster pulsar population. 2013. doi:10.1093/mnras/stt1850. 1309.7317.
- [62] C. Y. Hui, K. S. Cheng, and R. E. Taam. Dynamical Formation of Millisecond Pulsars in Globular Clusters. , **714** 1149, 2010. doi:10.1088/0004-637X/714/2/1149. 1003.4332.
- [63] R. N. Manchester, G. Fan, A. G. Lyne, V. M. Kaspi, and F. Crawford. Discovery of 14 Radio Pulsars in a Survey of the Magellanic Clouds. **649** 235, 2006. doi:10.1086/505461.
- [64] J. P. Ridley, F. Crawford, D. R. Lorimer, S. R. Bailey, J. H. Madden, R. Anella, and J. Chenamangalam. Eight new radio pulsars in the Large Magellanic Cloud. , **433** 138, 2013. doi:10.1093/mnras/stt709. 1304.6412.
- [65] R. N. Manchester, G. B. Hobbs, A. Teoh, and M. Hobbs. The Australia Telescope National Facility Pulsar Catalogue. **129** 1993, 2005.
- [66] P. Demorest, G. C. Bower, S. Chatterjee, J. Cordes, J. Deneva, J. Dexter, M. Kramer, T. J. W. Lazio, S. M. Ransom, L. Shao, N. Wex, and R. Wharton. ngVLA searches for pulsars at the Galactic center. In *American Astronomical Society Meeting Abstracts #233*, volume 233 of *American Astronomical Society Meeting Abstracts*, page 361.24. 2019.
- [67] V. M. Kaspi. The Neutron Star Zoo. In P. Weltevrede, B. B. P. Perera, L. L. Preston, and S. Sanidas, editors, *Pulsar Astrophysics the Next Fifty Years*, volume 337 of *IAU Symposium*, pages 3–8. 2018. doi:10.1017/S1743921317010390. 1806.03697.
- [68] R. A. Hulse and J. H. Taylor. A high sensitivity pulsar survey. **191** L59, 1974.
- [69] D. C. Backer, S. R. Kulkarni, C. Heiles, M. M. Davis, and W. M. Goss. A millisecond pulsar. **300** 615, 1982.
- [70] M. Burgay, N. D’Amico, A. Possenti, R. N. Manchester, A. G. Lyne, B. C. Joshi, M. A. McLaughlin, M. Kramer, J. M. Sarkissian, F. Camilo, V. Kalogera, C. Kim, and D. R. Lorimer. An increased estimate of the merger rate of double neutron stars from observations of a highly relativistic system. **426** 531, 2003.
- [71] A. G. Lyne, M. Burgay, M. Kramer, A. Possenti, R. N. Manchester, F. Camilo, M. A. McLaughlin, D. R. Lorimer, N. D’Amico, B. C. Joshi, J. Reynolds, and P. C. C. Freire. A double-pulsar system: A rare laboratory for relativistic gravity and plasma physics. **303** 1153, 2004.
- [72] F. Camilo, S. M. Ransom, J. P. Halpern, J. Reynolds, D. J. Helfand, N. Zimmerman, and J. Sarkissian. Transient pulsed radio emission from a magnetar. , **442** 892, 2006. arXiv:astro-ph/0605429.
- [73] D. R. Lorimer, M. Bailes, M. A. McLaughlin, D. J. Narkevic, and F. M. Crawford. A bright extragalactic radio burst. 2007.

- [74] M. Bailes, S. D. Bates, V. Bhallerao, N. D. R. Bhat, M. Burgay, S. Burke-Spolaor, N. D’Amico, S. Johnston, M. J. Keith, M. Kramer, S. R. Kulkarni, L. Levin, A. G. Lyne, S. Milia, A. Possenti, L. Spitler, B. Stappers, and W. van Straten. Transformation of a Star into a Planet in a Millisecond Pulsar Binary. *Science*, **333** 1717, 2011. doi:10.1126/science.1208890. 1108.5201.
- [75] D. L. Kaplan, K. Stovall, M. H. van Kerkwijk, C. Fremling, and A. G. Istrate. A Dense Companion to the Short-period Millisecond Pulsar Binary PSR J0636+5128. , **864** 15, 2018. doi:10.3847/1538-4357/aad54c. 1807.04610.
- [76] S. E. Thorsett, Z. Arzoumanian, and J. H. Taylor. PSR B1620–26: A binary radio pulsar with a planetary companion? **412** L33, 1993.
- [77] S. E. Thorsett, Z. Arzoumanian, F. Camilo, and A. G. Lyne. The Triple Pulsar System PSR B1620–26 in M4. **523** 763, 1999.
- [78] Z. Arzoumanian, K. Joshi, F. Rasio, and S. E. Thorsett. Orbital parameters of the PSR B1620–26 triple system. pages 525–530.
- [79] S. M. Ransom, I. H. Stairs, A. M. Archibald, J. W. T. Hessels, D. L. Kaplan, M. H. van Kerkwijk, J. Boyles, A. T. Deller, S. Chatterjee, A. Schechtman-Rook, A. Berndsen, R. S. Lynch, D. R. Lorimer, C. Karako-Argaman, V. M. Kaspi, V. I. Kondratiev, M. A. McLaughlin, J. van Leeuwen, R. Rosen, M. S. E. Roberts, and K. Stovall. A millisecond pulsar in a stellar triple system. , **505** 520, 2014. doi:10.1038/nature12917. 1401.0535.
- [80] A. M. Archibald, N. V. Gusinskaia, J. W. T. Hessels, A. T. Deller, D. L. Kaplan, D. R. Lorimer, R. S. Lynch, S. M. Ransom, and I. H. Stairs. Universality of free fall from the orbital motion of a pulsar in a stellar triple system. , **559** 73, 2018. doi:10.1038/s41586-018-0265-1. 1807.02059.
- [81] Y. Shao and X.-D. Li. Black hole/pulsar binaries in the Galaxy. , **477** L128, 2018. doi:10.1093/mnras/sly063. 1804.06014.
- [82] C.-A. Faucher-Giguère and A. Loeb. Pulsar-black hole binaries in the Galactic Centre. , **415** 3951, 2011. doi:10.1111/j.1365-2966.2011.19019.x. 1012.0573.
- [83] C. S. Ye, K. Kremer, S. Chatterjee, C. L. Rodriguez, and F. A. Rasio. Millisecond Pulsars and Black Holes in Globular Clusters. arXiv e-prints, 2019. 1902.05963.
- [84] M. Kramer, I. Stairs, R. Manchester, M. McLaughlin, A. Lyne, R. Ferdman, M. Burgay, D. Lorimer, A. Possenti, N. D’Amico, J. Sarkissian, G. Hobbs, J. Reynolds, P. Freire, and F. Camilo. Tests of general relativity from timing the double pulsar. *Science*, **314** 97, 2006. doi:10.1126/science.1132305. arXiv:astro-ph/0609417.
- [85] M. A. McLaughlin. The International Pulsar Timing Array: A Galactic Scale Gravitational Wave Observatory. arXiv e-prints, 2014. 1409.4579.